

HIV Medication Chart

Nucleoside/Nucleotide Analogue Reverse Transcriptase Inhibitors (NRTI)

Emtriva*
(emtricitabine, FTC)



Epivir*
(lamivudine, 3TC)



Retrovir*†
(zidovudine, AZT, ZDV)



Videx EC†
(didanosine, ddl)



Viread
(tenofovir, TDF)*



Zerit*
(stavudine, d4T)



Ziagen*
(abacavir, ABC)



Protease Inhibitors (PI)

Aptivus
(tipranavir, TPV)



Crixivan
(indinavir, IDV)



Invirase
(saquinavir hard gel capsules, SQV)



Kaletra*
(lopinavir/ritonavir, LPV/r)



Lexiva
(fosamprenavir, FPV)



Norvir*
(ritonavir, RTV)



Prezista
(darunavir, DRV)



Reyataz
(atazanavir, ATV)



Viracept
(nelfinavir, NFV)



Fixed Dose Combinations

Atripla
(TDF+FTC+EFV)



Combivir
(AZT plus 3TC)



FDA Pregnancy Category D


Epzicom
(ABC plus 3TC)



Trizivir
(AZT plus 3TC plus ABC)



Truvada
(TDF plus FTC)



Non-Nucleoside Reverse Transcriptase Inhibitors (NNRTI)

Intence
(etravirine, ETV)




Rescriptor
(delavirdine, DLV)



Sustiva*
(efavirenz, EFV)



Viramune*
(nevirapine, NVP)



FDA Pregnancy Category D

Entry Inhibitors

Fuzeon
(enfuvirtide, T-20)
Fusion Inhibitor



Selzentry
(maraviroc, MVC)
CCR5 Antagonist



Integrase Inhibitors

Isentress
(raltegravir, RAL)



All pills shown in actual size.

Medication brand names appear in bold. Generic names and commonly used abbreviations appear in parentheses.



* Also available in liquid form.

† Generic formulation available.

Medication Schedule

Name _____

Date _____

Name of Medication	Number of pills to take each time	Time of day you are to take this medicine		Food Interactions		Side Effects	TOTAL number of pills each day
				With Food <input type="checkbox"/>	Without Food <input type="checkbox"/>		

Helpful Hints:

- Refill prescriptions before you run out. Call for refills when you have at least 3 or 4 days left.
- Use cues as a reminder to take your pills (after a meal or favorite TV show, or before bedtime).
- Use reminder aids such as beepers, alarm clocks, and pillboxes. Ask your pharmacist about these.
- Plan ahead (vacations, travel, count out weekly doses).
- Do not stop taking your medications until you have spoken with your health care provider or pharmacist.
- If you have a severe reaction or in case of emergency, contact your health care provider IMMEDIATELY.

Contact Information: Provider _____

Clinic Phone _____ Pharmacy _____ Phone _____

Colorado AIDS Education and Training Center
University of Colorado Denver School of Medicine





UCD – A01 • 12631 E. 17th Avenue • PO Box 6511, MS A089 • Aurora, CO 80045

TEL: 303-724-0646 • FAX: 303-724-0875 • www.coloradoaetc.org

Developed by MeriLou Johnson, MSW, MPA and Steven Johnson, MD



Discontinued Medications or Formulations

<p>Agenerase (amprenavir, APV)</p> 	<p>Fortovase (saquinavir soft gel capsule, SQV)</p> 	<p>HIVID (zalcitabine, ddC)</p> 
<p>Kaletra (lopinavir/ritonavir, LPV/r)</p> 	<p>Videx (didanosine, ddl)</p> 